

Kinderkrankheiten

Erkrankung	Daten	Verlauf und Leitsymptome	Bilder
Keuchhusten	<ul style="list-style-type: none"> IfSG: §34 Bordetella pertussis toxinbildend Tröpfcheninfektion 7 - 14 Tage Impfkalender 4 x 1. Lj, alle 10J Gefahr: Lungenschäden, Enzephalopathie Säuglinge: Atemstillstand 	<p>The graph shows the course of pertussis. The y-axis represents temperature in °C, ranging from 37°C to 38°C. The x-axis represents time in days and weeks. The course is divided into four stages: IKZ (7-14 Tage), Stadium catarrhale (7-14 Tage), Stadium convulsivum (3-6 Wochen), and Rekonvaleszenz (mehrere Wochen). Symptoms include dry cough, sneezing, and catarrhal symptoms in the first stage; severe coughing fits with inspiratory stridor and cyanosis in the second stage; and convulsions in the third stage. The infection is contagious for up to 6 weeks.</p>	
Diphtherie	<ul style="list-style-type: none"> IfSG: §6, 7, 34 Corynebacterium diphtheriae Tröpfcheninfektion, Schmierinfektion 2 - 5 Tage Impfkalender 4 x 1. Lj, alle 10J Gefahr: Toxischer Verlauf, Myokarditis 	<p>The graph shows the course of diphtheria. The y-axis represents temperature in °C, ranging from 37°C to 40°C. The x-axis represents time in days. The course is divided into two stages: IKZ (2-5 Tage) and Organmanifestationsstadium (5-10 Tage). Symptoms include nasal diphtheria (bloody mucus), pharyngeal diphtheria (pseudomembrane), and laryngeal diphtheria (croup). The toxic form involves high fever, heart failure, and myocarditis.</p>	
Asthma	<ul style="list-style-type: none"> Anfallsweise Obstruktion untere Atemwege Allergisch IgE-vermittelt oder nicht-allergisch Beruhigen, Kutschersitz, Lippenbremse Sympathomimetika, Glukokortikoide 	<ul style="list-style-type: none"> Trockener Husten Hypersonor Pfeifen, Giemen, Brummen expiratorischer Stridor Erhöhter Atemwegswiderstand, Tiffenau-Test 	
Pseudokrupp	<ul style="list-style-type: none"> Akute subglottische Verengung Viral (Parainfluenzaviren) oder allergisch Kleinkinder, viele Rezidive Feuchte kalte Luft ist gut, Adrenalin, Glukokortikoide 	<ul style="list-style-type: none"> Heiserkeit Bellender Husten in- und expiratorischer Stridor selten Fieber meist Nachts Zyanose Starke Unruhe 	
Epiglottitis	<ul style="list-style-type: none"> NOTFALL!! Hoch-Akut bakteriell v.a. durch Haemophilus influenzae b Kleinkinder bis 5. Lj, Antibiotika Rachenuntersuchung nur in Intubationsbereitschaft 	<ul style="list-style-type: none"> Starke Schluckschmerzen kloßige Sprache inspiratorischer Stridor Todesangst Hypersalivation hohes Fieber 	